

The v_c - σ_c relation in low mass and low surface brightness galaxies

P. Buyle^{1,2,*†}, L. Ferrarese², G. Gentile³, H. Dejonghe¹, M. Baes¹, U. Klein⁴

¹*Sterrenkundig Observatorium, Universiteit Gent, Krijgslaan 281, S9, B-9000, Ghent, Belgium*

²*Herzberg Institute of Astrophysics, National Research Council of Canada, Victoria, BC V9E2E7, Canada*

³*University of New Mexico, Department of Physics and Astronomy, 800 Yale Blvd NE, Albuquerque, New Mexico 87131, USA*

⁴*Bonn University, Auf dem Hügel 71D, 53121 Bonn, Germany*

Accepted 1988 December 15. Received 1988 December 14; in original form 1988 October 11

ABSTRACT

We present an updated investigation of the relation between large scale disk circular velocity, v_c , and bulge velocity dispersion, σ_c . New bulge velocity dispersions are measured for a sample of 11 low surface brightness (LSB) and 7 high surface brightness (HSB) spiral galaxies for which v_c is known from published optical or HI rotation curves. We find that, while LSB galaxies appear to define the upper envelope of the region occupied by HSB galaxies (having relatively larger v_c for any given σ_c), the distinction between LSB and HSB galaxies in the $v_c - \sigma_c$ plane becomes less pronounced for $\sigma_c \lesssim 80 \text{ km s}^{-1}$. We conclude that either the scatter of the $v_c - \sigma_c$ relation is a function of v_c (and hence galaxy mass) or that the character of the $v_c - \sigma_c$ relation changes at $v_c \sim 80 \text{ km s}^{-1}$. Some implications of our findings are discussed.

Key words: black hole physics — dark matter — galaxies: haloes — galaxies: nuclei

1 INTRODUCTION

The existence of supermassive black holes (SBHs) in galactic nuclei has been accepted since the detections in the nearby spiral NGC 4258 (Miyoshi et al. 1995) and in our own galaxy (Schödel et al. 2003; Ghez et al. 2003). Compelling cases now exist in three dozen additional galaxies (see Ferrarese & Ford 2005 for a review), making it possible to investigate the existence of scaling relations linking the SBH mass, M_{BH} , to the overall properties of the host. Indeed, M_{BH} correlates with the blue luminosity L_B of the host bulge (Kormendy & Richstone 1995; Marconi & Hunt 2003), with the central light concentration (Graham et al. 2001), and with the bulge velocity dispersion σ_c (Ferrarese & Merritt 2000; Gebhardt et al. 2000). Because of its tight scatter (0.34 dex in M_{BH}), the $M_{BH} - \sigma_c$ relation is thought to allow us to peer into the mechanisms controlling the joint formation/evolution of SBHs and galaxies. However, cosmological simulations show that a possibly more fundamental relation should be expected between M_{BH} and the total gravitational mass M_{tot} of the galaxy, or the mass M_{DM} of the dark matter (DM) halo (e.g. Di Matteo et al. 2003; Kawakatu et al. 2005; Wyithe & Loeb 2005; Adams, Graff & Richstone

2000; Monaco, Salucci & Danese 2000; Haehnelt, Natarajan & Rees 1998; Cattaneo, Haehnelt & Rees 1999; Silk & Rees 1998; Haehnelt & Kauffmann 2000; Loeb & Rasio 1994). Indirect observational evidence for such a link has been found in the form of a correlation between the circular velocity v_c of spirals' disks (obtained by either HI or deep optical observations) and the σ_c of their bulge component (Ferrarese 2002; Baes et al. 2003; Pizzella et al. 2005):

$$\log v_c = (0.84 \pm 0.09) \log \sigma_c + (0.55 \pm 0.29) \quad (1)$$

Further exploring the $v_c - \sigma_c$ relation is of interest because, by linking galactic components on vastly different scales, the relation is a reflection of the interplay between disks and spheroids during galaxy formation/evolution. Furthermore, if v_c and σ_c are used as surrogates for M_{DM} and M_{BH} respectively, equation (1) implies a strong causal relation between SBHs and DM haloes.

Pizzella et al. (2005) report evidence of a separation in the $v_c - \sigma_c$ plane between High and Low Surface Brightness spiral galaxies (HSB and LSB respectively), the latter having distinctly larger v_c at a given σ_c compared to the former. Based on this, the authors argue against the importance of baryonic collapse in shaping the density profiles of DM haloes in LSBs, and suggest that for a given DM halo mass, LSBs might host less massive SBHs than HSBs. The behaviour of HSBs themselves is somewhat controversial: while Pizzella et al. (2005) find that HSB in the range

* E-mail: Pieter.Buyle@UGent.be

† Post-doctoral Fellow of the Fund for Scientific Research - Flanders, Belgium (F.W.O.)




Figure 1. Spectra for four of the program galaxies are shown in black. The top two panels show spectra obtained at the VLT, while the bottom two panels show Calar Alto spectra. To illustrate the quality of the data, for each telescope we chose to show the two spectra with the lowest (first and third panel from the top) and highest (second and fourth panel from the top) signal-to-noise ratio. The red lines show the best χ^2 fits to each spectrum, overplotted to the real data in the spectral region used to perform the fits. The narrow emission features present in three of the spectra are artifacts resulting from imperfect sky subtraction

$50 \lesssim \sigma_c \lesssim 350 \text{ km s}^{-1}$ (corresponding to $150 \lesssim v_c \lesssim 550 \text{ km s}^{-1}$) follow a linear relation, Ferrarese (2002) and Baes et al. (2003) claim that the relation is mildly non-linear for $\sigma_c \gtrsim 80 \text{ km s}^{-1}$ ($v_c \gtrsim 140 \text{ km s}^{-1}$), and breaks down altogether at lower σ_c , possibly revealing the inability of the least massive haloes to form central SBHs (Shankar et al. 2006; Haehnelt, Natarajan & Rees 1998; Silk & Rees 1998; Loeb & Rasio 1994)

The goal of this paper is to investigate the $v_c - \sigma_c$ relation for a combination of LSB and HSB galaxies, with v_c in the critical range $110 - 220 \text{ km s}^{-1}$. The data reduction and analysis are discussed in §2 and §3 respectively. Discussion and conclusions can be found in §4.

2 OBSERVATIONS AND DATA REDUCTION

To compare the behaviour of LSBs and HSBs in the $v_c \lesssim 220 \text{ km s}^{-1}$ range of the $v_c - \sigma_c$ relation, we drafted a sample of galaxies with v_c known from published rotation curves, and proceeded to obtain new optical spectra from which to measure σ_c . We targeted 11 LSB galaxies with optical rotation curves from the Palunas & Williams (2000) compilation, and seven HSB galaxies with HI rotation curves from the Ursa Major sample of Verheijen (2001). Besides the need of selecting galaxies visible in the given observing season, and bright




Figure 2. Velocity and velocity dispersion curves for ESO215-G39 (left panel) and ESO376-G10 (right panel) derived by the direct χ^2 method. The vertical dashed lines are drawn at the location of the bulge effective radius, r_e .

enough to produce spectra with the required signal-to-noise ratio (S/N), the one essential criterium in our sample selection is that the published rotation curves must be symmetric relative to the galaxy's centre, and reach an asymptotic value at large radii.

Details of the sample and observations are given in Table 1. The LSB galaxies were observed with the VLT/UT4 telescope of the European Southern Observatory on the nights of 2004 April 20, 29 and May 6. The holographic grism GRIS_1028z+29 on FORS2 was centred on the Calcium absorption triplet around $8500\text{\AA}$, producing an instrumental broadening $\sigma_{instr} \approx 35 \text{ km s}^{-1}$ with the $0''.7$ slit. The seven HSB galaxies were observed with the TWIN spectrograph at the 3.5m Calar Alto telescope, on 2005, June 6-19. The T06 grating was used in 1st order, again centred at the Ca triplet. The instrumental broadening was $\sigma_{instr} \approx 20 \text{ km s}^{-1}$ for the $1''.2$ wide slit. All spectra were divided into several exposures to ease cosmic ray identification and removal.

Standard data reduction was performed with MIDAS, IRAF and additional software developed specifically for this project. After basic data processing (dark and bias subtraction, flat fielding and trimming), atmospheric emission lines were identified and removed through an interpolation scheme, after which the spectra were rectified and wavelength calibrated. Cosmic rays were removed by applying a median filter along the spatial axis.

3 DATA ANALYSIS

Deriving accurate kinematical quantities of galaxies is a non-trivial task. The result of the extraction can vary depending on the applied fitting techniques, the S/N, the template stars, the absorption lines used, etc. (e.g., De Bruyne et al. 2003). To lower this degree of subjectivity, we derived velocity dispersions using two independent techniques: a direct χ^2 fit to the spectrum (De Bruyne et al. 2004), and the PPXF algorithm (Cappellari & Emsellem 2004) (see Fig. 1), which is based on a Penalized Likelihood approach. Velocity dispersions derived using these two methods agree to within 10%; the final values quoted in Table 1 were obtained using the χ^2




Figure 3. Left panel: The relation between the large scale disk circular velocity v_c and the central bulge velocity dispersion σ_c for HSBs (black circles) and LSBs (red triangles). Data from Ferrarese (2002), Baes et al. (2003), Pizzella et al. (2005) (full symbols) and this paper (open symbols) are plotted. Right panel: as in the left panel, with fits superimposed. The solid and dashed black lines are least square fits to HSB galaxies with $\sigma_c \geq 80 \text{ km s}^{-1}$ and $\sigma_c \leq 80 \text{ km s}^{-1}$ respectively. The red solid line is the fit to LSB galaxies with $\sigma_c \leq 80 \text{ km s}^{-1}$. Dotted lines represent the 1σ uncertainties in the fits.

routine. In analyzing the spectra, sky residuals that deviated by more than 3 times the average sigma in the nearby continuum were rejected; furthermore, to minimize the effects of template mismatch, six different template stars were used, each with weight calculated by a quadratic fit to the central row of the spectrum. In keeping with previous works, velocity dispersions were extracted from spectra binned, in the spatial direction, out to 1/8th of the bulge effective radius, estimated from the literature in the case of the HSB galaxies (Baggett et al. 1998; Courteau 1996; Heraudeau et al. 1996; Möllenhoff & Heidt 2001), and from published surface brightness profiles for the Palunas & Williams (2000) galaxies. Within this radius, the effect of a possible contamination by the disk rotational velocity should be negligible. This was verified by measuring the dispersion profiles along the major axis (without binning spatially); the velocity dispersion of most galaxies (see Fig. 2) indeed remained nearly constant in the bulge dominated region.

The circular rotational velocities and errors are published by Verheijen (2001) for the HSB galaxies. These were derived from HI observations and extend beyond the optical radius (R_{25}), well into the flat part of the curve. For the LSB galaxies, we measured v_c as the weighted average of the flat part of the optical rotation curve published in Palunas & Williams (2000); the error on v_c was derived by means of a bootstrap method and reflects any slight asymmetries in the rotation curves. The optical rotation curves are not as extended as the HI observations of the HSB galaxies, however all rotation curves reach an asymptotic value at large radii. As shown in Pizzella et al. (2005), when this condition is satisfied, v_c can be extracted reliably; therefore we do not expect any systematic biases in the v_c measurements

between our samples of HSBs and LSBs. The results of the dynamical analysis are given in Table 1.

4 DISCUSSION AND CONCLUSIONS

The $v_c - \sigma_c$ relation for HSB galaxies is plotted in the left panel of Fig. 3. The 11 LSB galaxies studied in this paper populate the region between $40 \leq \sigma_c \leq 80 \text{ km s}^{-1}$, where previously only three galaxies could be found. Furthermore, in the same region, we more than double the number of HSB galaxies, from three to eight. Pizzella et al. (2005) conclude that, relative to the $v_c - \sigma_c$ relation defined by HSB galaxies, LSB galaxies appear to have larger circular velocities, v_c , at any given value of σ_c . This conclusion is reaffirmed by our data. However, with the addition of the present sample, it has now become more evident that there is some overlap in the locations occupied by LSB and HSB galaxies in the $v_c - \sigma_c$ plane. In other words, LSB galaxies appear to occupy the *upper envelope* of the strip defined by HSB galaxies. Furthermore, at low velocity dispersions $30 \leq \sigma_c \leq 80 \text{ km s}^{-1}$, the $v_c - \sigma_c$ relation for both LSB and HSB galaxies appears to be shifted towards larger circular velocities relative to the relation defined by the (HSB) galaxies with larger velocity dispersion.

A least-square fit, taking into account the errors on both quantities (FITEXY; Press et al. 2002), to the sample of HSBs with $\sigma_c \geq 80 \text{ km s}^{-1}$ gives:

$$\log v_c = (0.68 \pm 0.04) \log \left(\frac{\sigma_c}{129 \text{ km s}^{-1}} \right) + (2.33 \pm 0.01), \quad (2)$$

with $\chi_r^2 = 2.58$. For $30 \leq \sigma_c \leq 80 \text{ km s}^{-1}$, on the other hand,

$$\log v_c = (1.33 \pm 0.86) \log \left(\frac{\sigma_c}{63 \text{ km s}^{-1}} \right) + (2.21 \pm 0.03), \quad (3)$$

with $\chi_r^2 = 0.23$ for HSB galaxies, and

$$\log v_c = (1.91 \pm 0.68) \log \left(\frac{\sigma_c}{58 \text{ km s}^{-1}} \right) + (2.20 \pm 0.02), \quad (4)$$

with $\chi_r^2 = 1.55$ for LSB galaxies.

Although the uncertainties are large, equations (3) and (4) are consistent with each other well within 1σ , but inconsistent with equation (2) (see also right panel of Figure 3).

Two interpretations are open at this point. The samples studied to date are far from complete or homogeneous, and it is conceivable that, as more data are collected, more galaxies will fill the gap between the LSB galaxies, which are mainly found in the low-velocity dispersion region, and the extrapolation to low σ_c of equation (2), defined mainly by HSB galaxies. This interpretation allows for the possibility that the log-log $v_c - \sigma_c$ relation, including both HSB and LSB galaxies, is linear, but that its *scatter* increases going from large to small velocity dispersions. Alternatively, it is possible that the log-log $v_c - \sigma_c$ relation is not linear, and that indeed its character changes dramatically below $v_c \sim 200 \text{ km s}^{-1}$, as originally suggested by Ferrarese (2002) and Baes et al. (2003). In either case, it seems unavoidable to conclude that the efficiency of bulge (and, perhaps, SBH) formation, is regulated by intrinsic parameters in addition to the depth of the global potential well.

The implications of these findings for the central SBHs cannot be easily quantified. The $M_{BH} - \sigma_c$ relation is not characterised for LSBs, for which no SBH detection has been attempted. It is also not defined below $\sigma_c \leq 80 \text{ km s}^{-1}$, which corresponds to SBHs too small to be detected given current space or ground-based instrumentation. We reiterate the conclusion of Ferrarese et al. (2002) and Baes et al. (2003) that the behaviour of the $v_c - \sigma_c$ relation at $\sigma_c \lesssim 80 \text{ km s}^{-1}$ might reflect the inability of these galaxies to form a central SBH, as argued on theoretical grounds by, for instance, Shankar et al. (2006); Haehnelt, Natarajan & Rees (1998); Silk & Rees (1998); Loeb & Rasio (1994). Numerical simulations (Di Matteo et al. 2003; Robertson et al. 2005) predict that the $M_{BH} - \sigma_c$ relation originates from the feedback between the central SBH and the progenitor of the hot stellar component at the early stages of the formation of both; therefore, in low-mass, late-type HSB and LSB galaxies internal factors might lead to the suppression of such feedback, creating a dynamical relation between disk, bulge and SBH which is starkly different from that displayed by more massive systems.

ACKNOWLEDGMENTS

We would like to thank S. De Rijcke for sharing his software to derive velocity dispersion profiles with us, P. Palunas for providing us the rotational velocities in tabular form and A. Pizzella for the very fruitful discussions. PB wishes to thank the Herzberg Institute of Astrophysics for the hospitality which made this investigation possible. Based on observations made at the European Southern Observatory, Chile (ESO Programme No. 073.B-0780). Based on observations collected at the Centro Astronómico Hispano Alemán (CAHA) at Calar Alto, operated jointly by the Max-Planck

Institut für Astronomie and the Instituto de Astrofísica de Andalucía (CSIC). PB acknowledges the Fund for Scientific Research Flanders (FWO) for financial support. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

REFERENCES

- Adams, F.C., Graff, D.S., Richstone, D., 2000, ApJ, 551, L31
- Baes, M., Buyle, P., Hau, G.K.T., Dejonghe, H., 2003, MNRAS, 341, L44
- Baggett, W. E., Baggett, S. M., Anderson, K. S. J., 1998, AJ, 116, 1626
- Cappellari, M. & Emsellem, E., 2004, PASP, 116, 138
- Cattaneo, A., Haehnelt, M.G., Rees, M.J., 1999, MNRAS, 308, 77
- Courteau, S., 1996, ApJS, 103, 363
- De Bruyne, V., Vauterin, P., De Rijcke, S., Dejonghe, H., 2003, MNRAS, 339, 215
- De Bruyne, V., De Rijcke, S., Dejonghe, H., Zeilinger, W.W. 2004, MNRAS, 349, 461
- De Rijcke, S., Michielsen, D., Buyle, P., Zeilinger, W.W., Dejonghe, H., Hau, G.K.T., 2005, AN, 326, 542
- Di Matteo, T., Croft, R. A.C., Springel, V., Hernquist, L. 2003, ApJ, 593, 56
- Ferrarese, L., 2002, ApJ, 578, 90
- Ferrarese, L. & Merritt, D., 2000, ApJ, 539, L9
- Ferrarese, L., & Ford, H.C. 2005, Sp.Sc.Reviews, 116, 523
- Gebhardt K. et al., 2000, ApJ, 539, L13
- Ghez, A.M., et al. 2003, ApJ, 586, L127
- Graham, A.W., Erwin, P., Caon, N., & Trujillo, I., 2001, ApJ, 563, L11
- Haehnelt, M.G., Natarajan, P., Rees, M.J., 1998, MNRAS, 300, 817
- Haehnelt, M.G., & Kauffmann, G., 2000, MNRAS, 318, L35
- Harmes, R.J., et al. 1995, ApJ, 435, L35
- Heraudeau, P., Simien, F., Mamon, G. A., 1996, A&A, 117, 417
- Hopkins, P. F., Hernquist, L., Martini, P., Cox, T. J., Robertson, B., Di Matteo, T., Springel, V., 2005, ApJ, 625, L71
- Kawakatu, N., Saitoh, T.R., Wada, K. 2005, ApJ, 628, 129
- Kelson, D., 2003, PASP, 115, 688
- Kormendy, J. & Richstone, D., 1995, ARA&A, 581
- Loeb, A. & Rasio, F., 1994, ApJ, 432, L52
- Marconi, A., & Hunt, L.K. 2003, ApJ, 589, L21
- Miyoshi, M., Moran, J., Herrnstein, J., Greenhill, L., Nakai, N., Diamond, P., & Inoue, M. 1995, Nature, 373, 127
- Möllenhoff, C., Heidt, J., 2001, A&A, 368, 16
- Monaco, P., Salucci, P., Danese, L., 2000, MNRAS, 311, 279
- Palunas, P. & Williams, T.B., 2000, AJ, 120, 2884
- Pizzella, A., Corsini, E.M., Dalla Bontá, E., Sarzi, M., Cocato, L., Bertola, F., 2005, ApJ, 631, 785
- Press, W.H., Teukolsky, S.A., Vetterling, W.T., Flannery,

Name	R.A. (J2000)	Dec. (J2000)	v_c (km s ⁻¹)	σ_c (km s ⁻¹)	r_e (arcsec)	v_{rad} (km s ⁻¹)	Morph	m_I (mag)	Incl. (degrees)	Exp.Time (seconds)	S/N (@ 8500 Å)
LSB galaxies											
ESO215-G39	11h17m04s	-49° 12' 05''	162±3	71±11	4.5	4335	SABc	12.01	50	500	20.03
ESO268-G44	12h48m42s	-45° 00' 29''	190±9	70±9	3.5	3477	Sb	12.22	62	450	25.68
ESO322-G19	12h29m07s	-40° 40' 24''	141±9	49±10	0	3100	SBc	12.64	79	900	22.19
ESO323-G42	12h55m01s	-40° 58' 20''	158±4	54±9	4	4203	Sc	11.53	69	450	30.32
ESO323-G73	13h04m02s	-38° 11' 57''	163±7	70±7	2	4929	Sbc	12.44	48	450	35.29
ESO374-G03	09h51m57s	-33° 04' 25''	146±3	40±6	6	2931	SABc	11.51	71	600	21.80
ESO382-G06	13h05m32s	-32° 57' 38''	171±4	54±8	2	4809	Sab	13.20	54	600	21.01
ESO444-G21	13h23m31s	-30° 06' 52''	116±21	53±9	4.5	4265	Sc	12.86	84	1900	20.46
ESO444-G47	13h28m25s	-31° 51' 28''	154±6	61±14	0	4389	SBc	12.82	71	900	22.01
ESO509-G91	13h40m03s	-25° 28' 28''	203±9	62±11	6.5	5113	SBc	12.70	79	900	20.91
ESO376-G10	10h42m00s	-36° 56' 07''	220±11	54±12	7	3184	SBd	11.00	76	600	20.27
HSB galaxies											
NGC 3953	11h53m49s	+52° 19' 36''	223±5	115±9	66	1052	Sbc	9.03	63	2500	60.01
NGC 3877	11h46m08s	+47° 29' 40''	167±11	84±8	53.5	895	Sc	9.91	83	3600	66.08
NGC 4088	12h05m34s	+50° 32' 21''	173±14	62±7	76	757	SABc	9.51	71	3600	58.13
NGC 3949	11h53m41s	+47° 51' 32''	164±7	63±5	36	800	Sbc	10.23	56	4000	57.15
NGC 4157	12h11m04s	+50° 29' 05''	185±10	71±6	35	774	SABb	9.93	90	4000	63.30
NGC 3769	11h37m44s	+47° 53' 35''	122±8	80±7	29	737	Sb	11.08	78	8000	66.25
NGC 3726	11h33m21s	+47° 01' 45''	162±9	63±10	84	866	Sc	9.50	49	6600	38.44

Table 1. Column 1 gives the galaxy's name, while columns 2 & 3 give the J2000 coordinates. The asymptotic circular velocity v_c is tabulated in column 4, it is taken directly from Verheijen (2001) for the HSB galaxies, and measured from the published rotation curves for the LSB galaxies. Central velocity dispersion (σ_c) at $r_e/8$ and the effective radius of the galaxy can be found in columns 5 and 6. The heliocentric radial velocity, Hubble classification, total I band magnitude, inclination can be found in columns 7 to 10 and are taken from Verheijen (2001) for the HSB galaxies and Palunas & Williams (2000) for the LSB galaxies and the online NED database. Finally, the total exposure time and S/N/pixel at 8500Å are given in columns 11 and 12.

- B.P., 2002, Numerical Recipes in C++. Cambridge Univ. Press, Cambridge
- Robertson, B., Hernquist, L., Cox, T.J., Di Matteo, T., Hopkins, P.F., Martini, P., Springel, V., 2005, astro-ph/0506038
- Schödel, R., Ott, T., Genzel, R., Eckart, A., Mouawad, N., & Alexander, T. 2003, ApJ, 596, 1015
- Shankar, F., Lap, A., Salucci, P., De Zotti, G., Danese, L. 2006, astro-ph/0601577
- Silk, J. & Rees, M.J., 1998, A&A, 331, L1
- Stuart, J. & Wyithe, B. 2006, MNRAS, 365, 1082
- Tremaine, S. et al., 2002, ApJ, 574, 740
- Van Zee, L., Skillman, E.D., Haynes, M.P., 2004, AJ, 128, 121
- Verheijen, M., 2001, ApJ, 563, 694
- Wyithe, J.S. & Loeb, A. 2005, ApJ, 634, 910

This paper has been typeset from a $\text{\LaTeX}$ file prepared by the author.